

# Eculizumab (Soliris)

Provider Order Form rev. 10/03/2024

## PATIENT INFORMATION

Referral Status:  New Referral  Updated Order  Order Renewal

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Patient Phone: \_\_\_\_\_  
Patient Address: \_\_\_\_\_ Patient Email: \_\_\_\_\_  
Allergies: \_\_\_\_\_  NKDA Weight (lbs/kg): \_\_\_\_\_ Height (in/cm): \_\_\_\_\_  
Sex:  M /  F Date of Last Infusion: \_\_\_\_\_ Next Due Date: \_\_\_\_\_ Preferred Location: \_\_\_\_\_

## DIAGNOSIS (Please provide ICD-10 code in space provided)

generalized myasthenia gravis without exacerbation: \_\_\_\_\_ Neuromyelitis Optica (NMOSD): \_\_\_\_\_  
Other: \_\_\_\_\_ Description: \_\_\_\_\_

## REQUIRED INFORMATION

MenACWY: Date of 1st dose: \_\_\_\_\_ Brand: \_\_\_\_\_  
Date of 2nd dose: \_\_\_\_\_ Brand: \_\_\_\_\_  
Meb B: Date of 1st dose: \_\_\_\_\_ Brand: \_\_\_\_\_  
Date of 2nd dose: \_\_\_\_\_ Brand: \_\_\_\_\_  
(Trumenba only) Date of 3rd dose: \_\_\_\_\_  
Prophylactic antibiotics prescribed:  Yes /  No  
Date patient started prophylactic antibiotics (if applicable): \_\_\_\_\_  
Provider REMS ID: \_\_\_\_\_

- For gMG diagnosis: Patient is anti-acetylcholine receptor antibody positive (provide documentation)
- For NMSOD diagnosis: Patient is anti-aquaporin-4 (AQP4) antibody positive (provide documentation)
- For gMG diagnosis: Meningococcal vaccine(s) given on \_\_\_\_\_ date. First Soliris dose may be given at least 2 weeks later unless otherwise specified.

## THERAPY ADMINISTRATION & DOSING (Choose one)

- Administer eculizumab (Soliris) 900mg weekly<sup>1</sup> x4 doses. Dilute with 90 ml 0.9% sodium chloride (final volume 180 ml) and infuse over 35 minutes.
  - Administer eculizumab (Soliris) 1200mg for the fifth dose one week after the fourth dose (week 5), then every 2 weeks<sup>1</sup> thereafter. Dilute with 120 ml 0.9% sodium chloride (final volume 240 ml) and infuse over 35 minutes.
  - If infusion is stopped for any reason, total infusion time should not exceed 2 hours
  - Monitor patient for hypersensitivity reaction for a period of 60 minutes following each infusion
- <sup>1</sup>Recommended dosage time intervals; may adjust +/- 2 days if needed

## LABORATORY ORDERS

Other: \_\_\_\_\_

## PRE-MEDICATION ORDERS

- Tylenol  500mg /  650mg PO
- Loratadine 10mg PO
- Pepcid 20mg  PO /  IVP
- Benadryl  25mg /  50mg  PO /  IVP
- Solumedrol  40mg /  125mg IVP
- Other: \_\_\_\_\_

## NURSING

- Hold infusion and notify provider for:
  - Signs/symptoms of infection or meningococcal infection such as:
    - Headache with (1) fever, (2) nausea/vomiting, (3) stiff neck/back
    - Muscle aches with flu-like symptoms, fever with or without rash, confusion or photophobia
- Ensure patient carries and understands Patient Safety Information Card.
- Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation.

## ADDITIONAL ORDERS

## PROVIDER INFORMATION

Preferred Contact Name: \_\_\_\_\_ Preferred Contact Email: \_\_\_\_\_  
Ordering Provider: \_\_\_\_\_ Provider NPI: \_\_\_\_\_  
Referring Practice Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_  
Practice Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

## REQUIRED DOCUMENTATION CHECKLIST (Additional documentation required for processing and insurance approval)

**Required Documentation:** Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications, Disease status, MRI, Flow Cytometry, MG classification, MG-ADL score, EMG results  
**Required Labs:** Anti-Ach receptor, Anti-AQP4,

Provider Name (print) \_\_\_\_\_ Provider Signature \_\_\_\_\_ Date \_\_\_\_\_