

Eculizumab (Soliris, BKEMV, Epysqli)



Provider Order Form rev. 06/23/2026

PATIENT INFORMATION

Referral Status: ☐ New Referral ☐ Updated Order ☐ Order Renewal

Patient Name:	DOB:	Patient Phone:
Patient Address:	Patient Email:	
Allergies:	☐ NKDA	Weight (lbs/kg):
Sex: ☐ M / ☐ F	Date of Last Infusion:	Height (in/cm):
Next Due Date:	Preferred Location:	

DIAGNOSIS (Please provide ICD-10 code in space provided)

Atypical Hemolytic Uremic Syndrome (aHUS):	Neuromyelitis Optica (NMOSD):
Paroxysmal Nocturnal Hemoglobinuria (PNH):	generalized Myasthenia Gravis (gMG):
Other:	Description:

REQUIRED INFORMATION

Men ACWY: Date of 1st dose: _____ Brand: _____
Date of 2nd dose: _____ Brand: _____

Men B: Date of 1st dose: _____ Brand: _____
Date of 2nd dose: _____ Brand: _____

(Trumenba only) Date of 3rd dose: _____

Prophylactic antibiotics prescribed: ☐ Yes / ☐ No

Date patient started prophylactic antibiotics (if applicable): _____

Provider REMS ID: _____

☐ For gMG diagnosis: Patient is anti-acetylcholine receptor antibody positive

☐ For NMOSD diagnosis: Patient is anti-aquaporin-4 (AQP4) antibody positive

THERAPY ADMINISTRATION & DOSING (Choose one)

☐ Administer Eculizumab or Eculizumab biosimilar.

☐ Administer this specific Eculizumab product: _____

aHUS, gMG, and NMOSD Diagnosis

☐ Administer 900mg weekly¹ x4 doses. Then administer 1200mg for the fifth dose one week after the fourth dose (week 5), then every 2 weeks¹ thereafter.

☐ Maintenance only: Administer 1200mg every 2 weeks¹.

PNH Diagnosis

☐ Administer 600mg weekly¹ x4 doses. Then administer 900mg for the fifth dose one week after the fourth dose (week 5), then every 2 weeks¹ thereafter.

☐ Maintenance only: Administer 900mg every 2 weeks¹.

¹Recommended dosage time intervals; may adjust +/- 2 days if needed

LABORATORY ORDERS

☐ Other: _____

PRE-MEDICATION ORDERS

☐ Tylenol ☐ 500mg / ☐ 650mg PO

☐ Loratadine 10mg PO

☐ Pepcid 20mg ☐ PO / ☐ IVP

☐ Benadryl ☐ 25mg / ☐ 50mg ☐ PO / ☐ IVP

☐ Solumedrol ☐ 40mg / ☐ 125mg IVP

☐ Other: _____

NURSING

☒ Hold infusion and notify provider for:

- Signs/symptoms of infection or meningococcal infection such as:
 - Headache with (1) fever, (2) nausea/vomiting, (3) stiff neck/back
 - Muscle aches with flu-like symptoms, fever with or without rash, confusion or photophobia

☒ Ensure patient carries and understands Patient Safety Information Card.

☒ Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation.

☒ If infusion is stopped for any reason, total infusion time should not exceed 2 hours

☒ Monitor patient for hypersensitivity reaction for a period of 60 minutes following each infusion

PROVIDER INFORMATION

Preferred Contact Name:	Preferred Contact Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

REQUIRED DOCUMENTATION CHECK LIST (Additional documentation required for processing and insurance approval)

Required Documentation: Patient demos, copy of front and back of primary and secondary insurance, 2 most recent OVN including treatment failures or contraindications, Disease status, MRI, Flow Cytometry, MG classification, MG-ADL score, EMG results

Required Labs: Anti-Ach receptor and/or Anti-AQP4

Provider Name (print)

Provider Signature

Date

Order valid for one year unless otherwise indicated. IV solutions/diluents may be substituted as allowed per manufacturer's instructions as necessitated by product availability.

Disclaimer: By signing this form, I authorize Novella Infusion and its affiliates to act as my designated agent in submitting prior authorizations, financial assistance applications, and other clinically required information with respect to this patient and order. This enrollment form shall serve as my signature for prior authorizations and financial assistance programs, as requested.